



Review Paper

Conventional and Molecular Improvement of Maize for Drought Tolerance: Review

Belay Fufa

Bako National maize research centre, Bako, Ethiopia

Corresponding author: belaygaroma@gmail.com

Article History: Received: June 10, 2024, Accepted: August 16, 2024

Abstract

Maize is an important staple crop for food, feed, bioenergy and industrial products globally. Despite the importance of maize as a principal food crop in developing countries, drought is a major constraint that affects maize production, particularly in Sub-Saharan Africa, where maize is grown under rainfed conditions. Plant breeders have been striving to improve and develop drought-tolerant variety. Nevertheless, these efforts still cannot meet the demand for food security due to fast population growth and climatic change. Conventional maize breeding for drought tolerance follows diverse approach includes recurrent selection, backcrossing, pedigree breeding, and subsequently evaluating inbred lines and hybrids at optimum conditions, managed screening site and random stress across multiple environments. Molecular markers were used to select donor parents with drought-adaptive alleles and then integrated into elite maize lines to create a new population of drought-tolerant maize inbred lines and subsequently used to develop hybrids maize tolerance to drought stress.

Keywords

Secondary traits, Molecular markers, Drought adaptive alleles, Maize

1. INTRODUCTION

Maize (*Zea mays* L.) is a worldwide important staple crop for food, feed, bioenergy and industrial products. Despite the importance of maize as a principal food crop, particularly in developing countries, its average yield in Africa (2.02 t ha^{-1}) is still low as compared to the world average (5.58 t ha^{-1}) (FAOSTAT, 2021). A significant portion of this yield gap is attributable to biotic and abiotic stresses. Adverse environments such as drought, salt, and high temperatures can affect maize growth and cause considerable yield loss. Among all, drought is a major constraint that affects crop production across the world, particularly in sub-Saharan Africa, where maize is grown under rainfed conditions. It is estimated about 15% to 20% of maize grain yield is lost each year due to drought, and even such losses may increase further because of climate change-induced rising temperatures and fluctuations in rainfall (Thomson et al., 2005; Cairns et al., 2013). Furthermore, climate change can further intensify the frequency of drought and significantly reduce maize grain yields (Mulungu and Ng'omb, 2019).

Drought affects at any maize growth stages. When drought occurred at the vegetative stage, it resulted in reduced leaf area, root spread, stem extension and metabolic activity. Also, it reduces the chlorophyll content and light interception, onward in low photosynthesis and subsequently

reduced maize yield (Athar and Ashraf, 2005). The fact that maize is more sensitive to drought at flowering or occurring between two weeks before and after the silking stage can cause significant yield loss of about 20% to 50% (Nielsen, 2007), Figure 1. If drought is severe and critical at flowering and extends throughout grain filling, it may lead to the complete abortion of ears, and the crop may become barren, and/or ears may have fewer kernels, which is very correlated with grain yield (Rhoads and Bennett, 1990; Ermeades, 2014).

So far, plant breeders have been striving to improve and develop drought-tolerant crops. Nevertheless, these efforts still cannot meet the demand of food security due to fast population growth and climatic change. Thus, the objective of this paper was to review the states of maize breeding for drought conditions using conventional breeding and molecular markers.

2. PLANTS RESPONSE TO DROUGHT STRESS

In an agricultural context, drought is a situation when the water availability of a plant is less than what is required to sustain its growth and development, or drought is a situation when there is inadequate moisture in the soil at a particular time to meet the needs of the plant. Plants use different mechanisms to cope with the drought. These strategies (mechanisms) include drought escape strategies,



Figure 1. Drought stress at different stages of maize development (Rhoads and Bennett, 1990).

i.e., early flowering time before the onset of the drought season, and maturation, meaning that plants undergo the completion of their life cycle prior to the detrimental effect of drought (Deikman et al., 2011). For example, Melkassa maize breeders in Ethiopia evaluated and selected maize germplasm that can escape drought via early flowering. So far, Melkassa1, Melkassa1Q, Melkassa4 (OPV), and hybrid maize (MH130, MH140, MH141) have been released and commercialized in low-moisture stress areas that receive annual rainfall < 800 mm in Ethiopia. Thus, farmers in low-moisture and marginal areas are using such varieties. However, a yield penalty can occur when too early varieties. Drought avoidance is also another strategy when plants improve water use efficiency through closing stomata, leaf rolling, increasing the thickness of the leaf cuticle, or improving water uptake via developing the root system. Plants also adapt metabolism mechanisms to minimize water loss, such as osmotic adjustment through active accumulation of solutes in the cell, which helps for retention of water and limits turgor loss (Paillard et al., 2003; Ribaut et al., 2009; Gowda et al., 2011). Traits like relative water, proline, and chlorophyll contents are also the most reliable parameters for drought tolerance in plants (Rahul et al., 2018). Drought escape and avoidance mechanisms are associated with morphological traits that can be exploited from diverse germplasm and selected at drought-prone areas.

3. FIELD MANAGEMENT FOR DROUGHT STRESS

Testing environments should be taken into consideration in relation to rainfall distribution, soil properties, altitude, genotype \times environment interaction and market demand. Evaluation of maize germplasm for drought tolerance should be carried out at optimum conditions under water stress during the rainy period using controlled irrigation, managed screening sites, and random stress. Water stress is in-

duced during the flowering and grain filling growth stages, and its average yield under stress could be reduced by up to 30% relative to the un-stressed yield (Maazou et al., 2020). Similarly, maize germplasm is induced at flowering time, which means drought stress coincides with anthesis and silk emergence, named severe stress (Banziger et al., 2000), and onwards supplementary irrigation should be applied.

3.1 Importance of Secondary Traits under Drought Tolerance in Maize Breeding

Grain yield under stressful conditions is often the primary and complex trait for selection. Secondary traits such as anthesis-silking interval, ear per plant, leaf senescence, stay green, chlorophyll content, and other morphological traits that are associated with grain yield should be considered under drought stress conditions (Lafitte et al., 2003; Banziger et al., 2000). Also, secondary traits should be genetically correlated with yield, exhibit genetic variation, have a high level of heritability, and be simple to measure (Ribaut et al., 2009). When drought coincided at anthesis silking interval, it affected embryonic development and early grain filling; yield loss estimated about 45% to 60% (Campos et al., 2006; Kumar et al., 2015). Furthermore, maize ovary abortion is greatest when drought is severed a few days before pollination, meaning that water deficiency around pollination increases the frequency of seed set abortion in maize due to a lack of photosynthate and to the distress of carbohydrate metabolism in ovaries (Zinselmeier et al., 1995). Banziger et al. (2000) reported that 90% decreased in yield as ASI increased from -0.4 to 10 days. On the other hand, the shorter ASI led to a positive association between ASI and grain yield and, indirectly, ear per plant and kernel per row selected. Mathieu and Aires. (2006) reported that ears per plant showed significant correlation with grain yield across drought ($r = 0.96$), low-N ($r = 0.66$) and optimum ($r = 0.83$) environmental conditions (Figure 2).

Delayed leaf senescence and high chlorophyll contents are also associated with the stay-green and key traits for drought tolerance. Stay green maize can retain the moisture and help better in photosynthesis during the grain filling stages (Lee and Tollenaar, 2007). Thus, plants survive with the existing soil moisture and maximize their grain filling efficiency. Consequently, breeders select genotypes that are capable of converting the sources to sink rapidly. For example, leaf growth and ASI are the secondary traits that determine the source and sink strengths of maize via their relationships with light interception and yield, respectively. In addition, staying green and root architecture are the most important secondary traits to impart drought tolerance and indirectly contribute to yield (Nepolean et al., 2013). Thus, indirect selection through secondary traits that are associated with grain yield can improve the efficiency of maize breeding under drought conditions. Therefore, secondary traits are highly significant and associated with grain yield

and should be considered for selection under drought stress in the maize breeding program.




Figure 2. Correlation of ears with grain yield under drought, optimum and low-N

4. BREEDING SCHEME

Maize breeding for drought tolerance is carried out by CGIAR; CIMMYT and IITA with the collaboration NARS in sub-Saharan Africa. At the beginning, intermediate and early-maturing stress-tolerant maize germplasm were tested in multiple countries in a wide range of environments and selected. Nearly two decades ago, the African Maize Stress (AMS) Project was launched with the collaboration IITA, CIMMYT and NARS in SSA to develop maize varieties and elite lines for drought tolerance (Menkir et al., 2014).

5. BROADENING GENE POOL

The first worth of a maize breeding program is broadening the gene pool, classifying inbred lines into heterotic groups using reliable opposite testers and markers, exploiting their variation among advanced inbreds, and subsequently assessing combining abilities among inbred lines for yield, drought tolerance, disease resistance, and other desirable agronomic traits. Broadening is step-wise through the introgression of drought-tolerant parents that carry favorable alleles or genes and recycle them for target traits. Breeding for drought tolerance is a state of motion to be continued to introduce exotic maize inbred lines from different genetic backgrounds that having novel alleles of adaptive drought-tolerant lines and then introgressed into elite lines to broaden germplasm and diversify the genetic base within and opposite heterotic groups. Maazou et al. (2020) reported that tropical inbred lines were characterized into

two heterotic groups using SNP markers, cluster analysis, testcross performance, and yield-based affinities. A similar study reported that heterotic grouping inbred lines fell into four groups, suggesting that each sub-group displayed the presence of substantial genetic diversity among these inbred lines, and this can maximize genetic variation for inbred line development and heterosis manifestation.

5.1 Conventional Breeding for Drought Tolerance in Maize

Maize breeding programs for drought tolerance follow diverse strategies such as recurrent selection, backcrossing, and pedigree breeding methods to exploit genetic variation while evaluating segregant population, fixed lines and hybrids under managed and across multi-environments at optimum and drought prone areas. The source of germplasm should be exhibited high allelic variation and diverse adapted parents that can promote drought tolerance Reynolds and Tuberosa (2008). The divers' sources include adapted elite lines developed from introgression, multi-parent populations and parental lines these carrying favorable alleles for drought tolerance that maximize genetic diversity within and opposite heterotic groups. Crossing within heterotic groups can start for inbred line development for target traits. For example: drought tolerance x drought tolerance for target traits; drought tolerance x heat tolerance, drought tolerance x low N tolerance, drought tolerance x disease resistance etc.

5.1.1 Performance of Maize Inbred Lines and Hybrids under Managed and Drought Stress

The aim of maize breeding for drought tolerance is 1) to reduce the yield gap between optimum and stress conditions, 2) to advance the yield stability across stress conditions. This can be employed by evaluating diverse maize germplasm under drought stress and optimum conditions, indicating that several inbred lines populations and hybrids screened under managed drought tolerance and optimum at the flowering and grain filling stages. Several studies have reported on the evaluation of inbred lines for drought tolerance. Chandel (2012) assessed maize inbred lines for ASI, plant height, leaf senescence, leaf rolling, and yield under limiting irrigation verses normal irrigation at knee height, flowering and grain filling. The performed inbred lines or families were selected based on high grain yields, short ASI, increased ears per plant, and reduced leaf senescence under managed drought stress were also assessed under watered conditions. Likewise, hundred maize inbred lines evaluated under contrasting soil moisture conditions, and among these fifteen inbred lines showed high tolerance to drought. Moreover, the tolerance inbred lines showed shorter ASI, a low canopy temperature, higher chlorophyll content and a comparatively higher grain yield when subjected to drought stress (Shadakshari and Shanthakumar, 2015). Likewise, eleven inbred lines evaluated under drought and heat over two years, and among these

two inbred lines revealed that they had a relatively high-water content and improved damage to reproductive tissues under drought conditions (Chen et al., 2012). Also, a high heritability was found for plant height, thousand kernel weight, grain rows per ear and grain yield per plant, in addition, the selected inbred lines showed high specific combining ability for yield under stress conditions. Several studies showed that physiological traits such as stomata conductance, chlorophyll content, transpiration rate, leaf water potential, and relative water content were used to improve breeding stress tolerance. For example, Dordas et al. (2018) reported a better association between physiological characteristics and grain yield under stress conditions for inbred lines than hybrids, and these traits are strongly helpful for the selection of adaptive parents. Similarly, a correlation of secondary traits with grain yield under stress conditions was conducted. Canopy temperature depression ($r = 0.61$) and chlorophyll present in sampled leaves ($r = 0.50$) showed positive correlation, while leaf rolling score ($r = -0.49^{**}$), leaf senescence score ($r = -0.57$) and anthesis silking interval ($r = -0.15$) showed negative correlation (Parajuli et al., 2018). This means that increasing leaf rolling and subsequent leaf drying early and non-nicking of ASI result in decreasing grain yield.

After the evaluation of several inbred lines, promising lines were selected based on their tolerance. The selected F4 stages should be crossed at least by two opposite testers, and their crosses should be evaluated under managed drought tolerance for direct selection and evaluated in a broad array of environments in drought-prone areas. The hybrid maize that was selected under stress might be reduced to at least 50% of its yield potential (i.e., if yield under optimum conditions is around 7 t ha^{-1} , yield under stress is likely about 3.5 t ha^{-1}) (Banziger et al., 2000). Forty drought-tolerance hybrids evaluated under well-watered drought and heat over three years and found a significant difference among hybrids for grain yield, and the yield showed a positive correlation with secondary traits (Meseka et al., 2018), and three hybrids produced a high grain yield under three conditions. Similarly, forty-nine maize hybrids that were developed via Line by test cross were evaluated under drought stress, low nitrogen, and optimum conditions and showed that the drought stress reduced grain yield and plant height. The authors also noticed that further maize improvement for grain yield is required under stress and optimum conditions (Ertiro et al., 2017).

5.2 Molecular Markers to Dissect Drought Tolerance Traits

Advanced genomics-assisted breeding techniques such as marker-assisted selection (MAS), marker-assisted backcrossing, and marker-assisted recurrent selection (MARS) are fast approaches and possibilities for drought tolerance improvement. Understanding the genetic and physiological basis of target traits is useful information during selection for drought tolerance. Quantitative trait locus (Quantitative

traits locus, QTL) is included both phenotypic data (trait measurement) and genotype data using DNA markers that attempt to uncover the genetic basis of variation in traits (Chamarthi et al., 2011).

5.2.1 Identification of Quantitative Traits Locus Associated with Drought Tolerance Traits

Drought tolerance is a complex nature in quantitative trait that is controlled by several genes and affects genotype by environment interaction. Many QTLs that control morpho-physiological traits and grain yield under water-limited conditions have been identified in maize. For instance, several QTLs detected across six chromosomes for secondary traits such as ASI, ear per plant, stay green and plant height (Almeida et al., 2014), Figure 3. Out of the detected QTLs, 65% were identified under water stress, suggesting the significance of secondary traits for maize breeding under drought tolerance. The authors also reported that twenty-nine QTL were detected for ear per plant and 80 QTL for stay green (leaf senescence and chlorophyll content) in maize populations under both conditions. Also, Zhao et al. (2018) found 62 QTLs for major agronomic traits, and out of these, nine QTLs were detected for ASI and ear length. Out of these, 75% (52 QTLs) were detected under water-stressed conditions, suggesting a high possibility for maize improvement under stress conditions.

It was interesting that clustered QTLs were found on chromosome 3 that harbored QTL (hotspot region) for most morpho-physiological traits (Almeida et al., 2014). Similarly, co-located QTLs were found on chromosomes 1, 3, and 5 for ear height and ASI under both conditions (Yang et al., 2008; Ribaut et al., 2009). It is suggesting that the same gene for controlling different traits or loci has a pleiotropic effect that controls multiple traits. Similarly, Zhao et al. (2018) also identified 36 meta-QTLs across 26 populations under stress and watered conditions and several candidate genes were identified through fine mapping, suggesting that these inbred lines can use as introgression into elite lines for drought tolerance. Additionally, a cluster of QTLs was detected for plant height, TKW, and grain yield per plant, and these may be used to enhance maize drought tolerance under water-limited environments (Chen et al., 2012).

The above results may suggest that molecular markers linked to secondary and desirable agronomic traits under stress enable the transfer of the donor parent into elite lines (Ribaut, 2002). Furthermore, molecular markers were used to select donor parents with drought-adaptive alleles and then integrated into elite maize lines to create a new population of drought-tolerant inbred lines. For example, Almeida et al. (2013) detected QTL in relation to GY with ASI and then introgressed into a drought-prone maize line, showing increased grain and lower ASI. Also, Bankole et al. (2017) reported that marker-assisted recurrent selection (MARS) helps to improve drought tolerance in population and yield

gain per cycle and increase the frequencies of favorable alleles.




Figure 3. QTLs detected across chromosomes for ASI, ear per plant, stay green, plant height ear ratio under stress and water regime (Wang et al., 2012; Almeida et al., 2014)

6. CONCLUSION

Drought is a major constraint that affects maize production, particularly in Sub-Saharan Africa, where maize is grown under rainfed conditions. Plant breeders have been striving to improve and develop drought-tolerant variety through conventional and molecular breeding approach. The conventional method includes recurrent selection, backcrossing and pedigree breeding meanwhile considering secondary traits for selection under drought stress and optimum that are directly associated with grain yield. Furthermore, several QTLs regulating important morpho-physiological traits were identified, molecular markers developed, and used to select donor parents with drought-adaptive alleles, and then integrated into elite maize lines to create a new source population of drought-tolerant inbred lines and subsequently develop hybrid resilience to drought stress.

ACKNOWLEDGMENT

-

REFERENCES

Almeida, G., S. M. C. Makumbi, A. Nair, J. B. JM, and R. B (2013). QTL mapping in three tropical maize populations reveals a set of constitutive and adaptive genomic regions for drought tolerance. *Theor Appl Genet*, (126); 583–600

Almeida, G., S. Nair, J. C. Borém, S. Trachsel, J. Ribaut, M. Bänziger, B. Prasanna, J. Crossa, and R. Babu (2014). Molecular mapping across three populations reveals a QTL hotspot region on chromosome 3 for secondary traits associated with drought tolerance in tropical maize. *Molecular Breeding*, (34); 701–715. <https://doi.org/10.1007/s11032-014-0068-5>

Athar, H. and M. Ashraf (2005). *Photosynthesis under drought stress*. Handbook of Photosynthesis (ed. M. Pessarakli)

Bankole, F., A. Menkir, G. Olaoye, J. Crossa, S. Hearne, N. Unachukwu, and M. Gedil (2017). Genetic gains in grain yield and other traits in a maize population improved using marker assisted recurrent selection. *Frontiers in Plant Science*, 8; 808

Banziger, M., O. Edmeades, D. Beck, and M. Bellon (2000). *Breeding for Drought and Nitrogen Stress Tolerance in Maize: from Theory to Practice*. CIMMYT, Mexico, DF)

Cairns, J., J. Hellin, K. Sonder, J. Araus, J. MacRobert, C. Thierfelder, and P. Prasanna (2013). Adapting maize production to climate change in sub-Saharan Africa. *Food Security*, 5; 345–360

Campos, H., M. Cooper, G. Edmeades, C. Loffer, J. Schussler, and M. Ibanez (2006). Changes in drought tolerance in maize associated with fifty years of breeding for yield in the US Corn Belt. *Maydica*, (51); 76–86

Chamarthi, S., A. Kumar, T. Vuong, M. Blair, P. Gaur, H. Nguyen, and R. Varshney (2011). *Trait mapping and molecular Breeding in legumes: concepts and examples in soybean, common bean and chickpea*. CABI International, Oxfordshire, UK (in press)

Chandel, U. (2012). Morphological traits determining drought tolerance in local and cimmyt maize inbreds (*Zea Mays L.*). *Agricultural Science Digest-A Research Journal*, 32(2); 160–163

Chen, J., W. Xu, J. Velten, Z. Xin, and J. Stout (2012). Characterization of maize inbred lines for drought and heat tolerance. *Journal of Soil and Water Conservation*, 67(5); 354–364

Deikman, J., M. Petracek, and J. Heard (2011). Drought tolerance through biotechnology: improving translation from the laboratory to farmers' fields. *Current Opinion in Biotechnology*, (23); 1–8

Dordas, C., F. Papathanasiou, A. Lithourgidis, J. Petrevska, I. Papadopoulos, C. Pankou, F. Gekas, E. Ninou, I. Mylonas, I. Sistanis, and C. Tzantarmas (2018). Evaluation of physiological characteristics as selection criteria for drought tolerance in maize inbred lines and their hybrids. *Maydica*, 63(2); 14

Ermeades, G. (2014). *Progress in Achieving and Delivering Drought Tolerance in Maize: An Update*. ISAAA, Ithaca

Ertiro, B., Y. Beyene, B. Das, S. Mugo, S. O. M. Olsen, C. Juma, M. Labuschagne, and B. Prasanna (2017). Combining ability and testcross performance of drought-tolerant maize inbred lines under stress and non-stress environments in Kenya. *Plant Breeding*, (136); 197–205

FAOSTAT (2021). *Statistical data bases and data-sets of the food and agricultural organization of the United nations*. <http://www.faostat.fao.org/default.aspx>

Gowda, V. R. P., A. Henry, A. Yamauchi, H. Shashidhar, and R. Serraj (2011). Root biology and genetic improvement for drought avoidance in rice. *Field Crops Research*, (122); 1–13

- Kumar, M., M. Uniyal, N. Kumar, S. Kumar, and R. Gangwar (2015). Conventional and molecular breeding for development of drought tolerant maize cultivars. *Journal Crop Sci Tech*, **4**; 1–3
- Lafitte, H., A. Blum, and G. Atlin (2003). *Adaptation strategies for agricultural water management under climate change in Europe*. International Rice Research Institute, Manila
- Lee, E. and M. Tollenaar (2007). Physiological basis of successful breeding strategies for maize grain yield. *Crop Science*, **4**; 202–215
- Maazou, A., J. Tu, J. Qiu, and Z. Liu (2020). Breeding for Drought Tolerance in Maize (*Zea mays* L.). *American Journal of Plant Sciences*, **7**; 1858–1870
- Mathieu, J. A. and F. Aires. (2006). Drought tolerance improvement in tropical maize source populations: evidence of progress. *Crop Sci*, **46**; 180–190
- Menkir, A., I. Dieng, M. Gedil, W. Mengesha, M. Oyekunle, P. Riberio, G. Adu, A. Yacoubou, M. Coulibaly, F. Bankole, and J. Derera (2014). Approaches and progress in breeding drought-tolerant maize hybrids for tropical lowlands in west and central Africa. Technical report, The Plant Genome
- Meseka, S., A. Menkir, B. Bossey, and W. Mengesha (2018). Performance Assessment of Drought Tolerant Maize Hybrids under Combined Drought and Heat Stress. *Scientific Reports*, **8**(12); 274
- Mulungu, K. and J. Ng'omb (2019). *Climate change impacts on sustainable maize production in sub-Saharan Africa*. A review. In A. Hossain (Ed.), *Maize—Production and use*
- Nepolean, T., I. Singh, F. Hossain, N. Pandey, and H. Gupta (2013). Molecular characterization and assessment of genetic diversity of inbred lines showing variability for drought tolerance in maize. *J. Plant Biochem Biotechnol*, **22**; 71–79
- Nielsen, R. (2007). *Assessing effects of drought on corn grain yield*. West Lafayette, In: *Purdue University*. <http://www.kingcorn.org/news/articles.07/Drought-0705.html>
- Paillard, S., T. Schnurbush, M. Winzeler, M. Messmer, P. Sourdille, O. Abderhalden, B. Keller, and G. Schachermayr (2003). An integrative genetic linkage map of winter wheat (*Triticum aestivum* L.). *Theoretical application Genetics*, **170**; 2176–2191
- Parajuli, S., B. Ojha, and G. Ferrara (2018). Quantification of secondary traits for drought and low nitrogen stress tolerance in inbreds and hybrids of maize (*Zea mays* L.). *J. Plant Genet. Breed*, **2**(1); 106
- Rahul, V., R. Panda, D. Lenka, and G. R. Rout (2018). Physiological and Biochemical Evaluation of Maize Hybrid Germplasm Lines for Drought Tolerance under Receding Soil Moisture Condition. *Int J Curr Microbiol App Sci*, **7**; 2176–2191
- Reynolds, M. and R. Tuberosa (2008). Translational research impacting on crop productivity in drought-prone environments. *Curr Opin Plant Biol*, **11**; 171–179
- Rhoads, F. and J. Bennett (1990). *Corn*. In: Stewart BA, Nielsen DR (eds) *Irrigation of agricultural crops*. ASA-CSSA-SSSA, Madison
- Ribaut, J. (2002). Use of molecular markers in plant breeding: drought tolerance improvement in tropical maize. In *Quantitative Genetics, Genomics, and Plant Breeding* (Kang, M.S., ed.). Technical report
- Ribaut, J., J. Betran, P. Monneveux, and T. Setter (2009). *Drought Tolerance in Maize*. In: Bennetzen, J. and Hake, S., Eds., *Handbook of Maize: Its Biology*. Springer, New York
- Shadakshari, T. and G. Shanthakumar (2015). Evaluation of maize inbred lines for drought tolerance under contrasting soil moisture regimes. In *Karnataka J. Agric. Sci.*, volume 28
- Thomson, A., N. Brown, R. Roseberg, C. Lazurade, and V. Benson (2005). Climate change impacts for the conterminous USA. An intergraded assessment, part 3. Dry land production of grains and forage crops. *Climate change*, **69**; 43–65
- Wang, A., Y. Li, and C. Zhang (2012). QTL Mapping for stay-green in maize (*Zea mays*). *Can J Plant Sci*, **92**; 249–256
- Yang, X., S. Lu, F. Zhang, Y. Zhou, Qu, and Cx.Xie (2008). QTL mapping of plant height and ear position in maize (*Zea mays* L.). *Hereditas*, **30**; 1477–1486
- Zhao, X., Y. Peng, J. Zhang, P. Fang, and B. Wu (2018). Identification of QTLs and Meta-QTLs for Seven Agronomic Traits in Multiple Maize Populations under Well-Watered and Water-Stressed Conditions. *Crop Science*, **58**; 507–520
- Zinselmeier, C., M. Westgate, J. Schussler, and R. Jones (1995). Low water potential disrupts carbohydrate metabolism in maize (*Zea mays* L.) ovaries. *Plant Physiol*, **107**; 385–391